

Question	Answers	Extra information	Mark	AO / Specification reference
01.1	zinc and copper chloride solution		1	AO2 4.4.1.2
01.2	hydrogen		1	AO1 4.1.2.5
01.3	less vigorously		1	AO1 4.1.2.5
01.4	lithium chloride		1	AO2 4.4.1.2
01.5	sodium > lithium > zinc > copper	one mark for one correct two marks for two correct three marks for all correct	3	AO2 4.4.1.2
02.1	reduced loses		1 1	AO1 4.4.1.3
02.2	$2\text{Fe}_2\text{O}_3 + 3\text{C} \rightarrow 4\text{Fe} + 3\text{CO}_2$		1	AO2 4.4.1.3
02.3	gold is unreactive so it is found as elemental in the earth		1 1	AO1 4.4.1.3
03.1	steady bubbling from surface of zinc <b>or</b> zinc gets smaller <b>or</b> clear gas is formed		1	AO1 4.4.1.2

Question	Answers	Extra information	Mark	AO / Specification reference
03.2	zinc chloride and hydrogen		1	AO2 4.4.1.2
03.3	$\text{Zn(s)} + 2\text{HCl(aq)} \rightarrow \text{ZnCl}_2\text{(aq)} + \text{H}_2\text{(g)}$	one mark for formulae and state symbols of reactants & one mark for formulae and state symbols of reactants <b>or</b> one mark for correct formulae & one mark for state symbols one mark for balancing	3	AO2 4.4.1.2 4.3.1.1
03.4	oxidised as it loses electrons/forms positive ions		1 1	AO2
04.1	Mg or Al does not react with $\text{CaCl}_2$ , so is below Ca in the reactivity series. reacts with copper sulfate and zinc chloride, so is above these two metals in the reactivity series.		1 1 1	AO2 × 2 AO3 × 1 4.4.1.2
04.2	$\text{X(s)} + \text{CuSO}_4\text{(aq)} \rightarrow \text{Cu(s)} + \text{XSO}_4\text{(aq)}$	one mark for formulae and state symbols of reactants & one mark for formulae and state symbols of reactants <b>or</b> one mark for correct formulae & one mark for state symbols one mark for balancing	3	AO2 4.4.1.2 4.3.1.1

Question	Answers	Extra information	Mark	AO / Specification reference
04.3	test tube full of aluminium salt solution add metal X if no observation, then metal is aluminium if displacement reaction occurs, metal is magnesium		1 1 1 1	AO2 4.4.1.2
05.1	mass of lead sulfide in the ore = $\frac{25}{100} \times 240 \text{ kg} = 60 \text{ kg}$ mass of one mole of PbS = $207 + 32 = 239 \text{ g}$ mass of Pb in 60 kg of PbS = $\frac{207}{239} \times 60 \text{ kg} = 51.96\dots$ = 52 kg		1 1 1 1	AO2 4.4.1.3
05.2	$2\text{PbS}(\text{s}) + 3\text{O}_2(\text{g}) \rightarrow 2\text{PbO}(\text{s}) + 2\text{SO}_2(\text{g})$	one mark for formulae and state symbols of reactants & one mark for formulae and state symbols of reactants <b>or</b> one mark for correct formulae & one mark for state symbols one mark for balancing	3	AO2 4.3.1.1
05.3	reduced because oxygen is lost/electrons are gained		1 1	AO1 4.4.1.3
05.4	lead is below carbon in the reactivity series but aluminium is above carbon in the reactivity series		1 1	AO1 4.4.1.3
05.5	<b>one</b> from: <ul style="list-style-type: none"> <li>• iron</li> <li>• copper</li> </ul>		1	AO1 4.4.1.3

Question	Answers	Extra information	Mark	AO / Specification reference
06	<b>Level 3:</b> A detailed and coherent comparison is given, demonstrating a sound knowledge and understanding of displacement reactions in Halogens and metals.		5-6	AO1 4.4.1.2 4.1.2.6
	<b>Level 2:</b> Correct descriptions of displacements are given for the Halogens and in metals. Some comparisons are made, but not all are clearly articulated.		3-4	
	<b>Level 1:</b> Some correct points are made about displacement reactions. Few comparisons are made, and these are not clearly articulated.		1-2	
	<b>No relevant content</b>		0	
	<b>Indicative content</b>			
	<ul style="list-style-type: none"> <li>in both the Halogen and metal reactions, a more reactive element displaces a less reactive element from a solution of its salt</li> <li>in both cases, one of the products is the less reactive Halogen or metal</li> <li>in both cases, the more reactive element ends up in a compound</li> <li>in the Halogen reaction, the more reactive Halogen is reduced (its atoms each gain one electron)</li> <li>in the metal reaction, the more reactive metal is oxidised (its atoms lose electrons)</li> <li>in the Halogen reaction, reactivity is based on tendency to form negative ions</li> <li>the closer the outer electron is to the nucleus, the more reactive the Halogen</li> <li>in the metal reaction, reactivity is based on the tendency to form positive ions</li> </ul>			

Question	Answers	Extra information	Mark	AO / Specification reference
07.1	lithium		1	AO2 4.4.1.2
07.2	magnesium oxide		1	AO2 4.4.1.1
07.3	oxidised magnesium has gained oxygen/lost electrons		1 1	AO1 AO2 4.4.1.1
07.4	magnesium is more reactive than carbon therefore, carbon cannot displace magnesium from magnesium oxide		1 1	AO1 4.4.1.3
08.1	nickel reacts more vigorously than lead with HCl, so it is above lead in the reactivity series nickel does not react with water, but iron does, so nickel is below iron in the reactivity series nickel is between iron and lead in the reactivity series		1 1 1	AO3 4.4.1.2
08.2	add nickel to a solution such as iron chloride there should be no reaction because nickel is less reactive than iron add iron to a solution of nickel chloride there should be a reaction because iron is more reactive than nickel		1 1 1 1	AO1 4.4.1.2
08.3	$\text{Ni} + \text{Cu}^{2+} \rightarrow \text{Ni}^{2+} + \text{Cu}$  copper	one mark for reactants, one mark for products	2  1	AO2 4.4.1.4
09.1	do not mix a metal with a solution containing the same metal	allow suitable examples (e.g. do not mix magnesium with magnesium chloride solution)	1	AO3 4.4.1.2

Question	Answers	Extra information	Mark	AO / Specification reference
09.2	copper of the three metals in the experiment, it is the lowest in the reactivity series <b>or</b> cannot displace any of the other metals		1 1	AO2 4.4.1.2
09.3	$\text{Mg(s)} + \text{Cu}^{2+}(\text{aq}) \rightarrow \text{Mg}^{2+}(\text{aq}) + \text{Cu(s)}$	one mark for formulae and state symbols of reactants & one mark for formulae and state symbols of reactants <b>or</b> one mark for correct formulae & one mark for state symbols one mark for balancing	3	AO2 4.4.1.4
09.4	before: 2,8,2 after: 2,8		1 1	AO2 4.4.1.7
09.5	reduces (potential harmful) waste after experiment cheaper, as smaller amounts of metals and solutions are required easier to observe on a white tile than in a test tube		1 1 1	AO3
10.1	there are no gaps/sticks between the nitrogen and hydrogen atoms		1	AO1 4.2.1.4
10.2	N atom should have 1 shell with 5 dots and 3 crosses. H atoms should have 1 shell with 1 dot and 1 cross each. H atoms should each be sharing 1 dot and 1 cross.		2	AO1 4.2.1.4
10.3	the molecules do not have an overall electric charge/no charged particles that are free to move		1	AO3 4.2.2.1

Question	Answers	Extra information	Mark	AO / Specification reference
10.4	gas		1	AO3 4.2.2.1
10.5	hydrazine has bigger molecules and the intermolecular forces between bigger molecules are stronger		1 1	AO1 4.2.2.4
11.1	metal atoms lose outer shell/highest energy level electrons in their reactions as atoms become bigger, their outer electrons are further from the nucleus (or more shielding) so the electrostatic attraction between the nucleus and outer electrons gets weaker so less energy is required to remove an electron		1 1 1 1	AO1 × 3 AO2 × 1 4.1.2.5 4.4.1.2
11.2	overall, radius decreases from top to bottom of the reactivity series this indicates that the conclusion is true calcium and magnesium do not fit the pattern, with both atoms being bigger than the one/two immediately above them in the reactivity series this indicates that the conclusion is not entirely correct overall, the trend identified in the conclusion is correct and the reason is correct		1 1 1 1 1	AO3
12.1	heat in a fume cupboard avoid handling the lead/wash hands		2	AO2
12.2	lead and carbon dioxide	both products required for the mark	1 1	AO2 4.4.1.2



Question	Answers	Extra information	Mark	AO / Specification reference
12.3	carbon dioxide has formed as a gas (and escaped from the evaporating basin)		1	AO1 4.3.1.3