

# Oxford Revise | Edexcel GCSE Maths Higher | Answers

## Chapter 29 Tables, averages, and range

Question	Answer	Extra information	Marks																												
29.1	$63 \times 10 = 630$ , the total of 10 of the numbers $51 \times 4 = 204$ , the total of 4 of the numbers $630 - 204 = 426$ , the total of the other six numbers The mean of these six numbers is $426 \div 6 = 71$	630 or 204 Subtracting the total of the four numbers from the total of the ten numbers. Correct answer	1  1  1																												
29.2 (a)	$\frac{71+1}{2} = 36$ The median is the 36th value This is in the class $30 < t \leq 35$	$\frac{71+1}{2} = 36$ $30 < t \leq 35$	1  1																												
29.2 (b)	<table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th>Time (<i>t</i> minutes)</th> <th>Frequency</th> <th>Midpoint</th> <th>Frequency × midpoint</th> </tr> </thead> <tbody> <tr> <td><math>20 &lt; t \leq 25</math></td> <td>10</td> <td>22.5</td> <td>225</td> </tr> <tr> <td><math>25 &lt; t \leq 30</math></td> <td>17</td> <td>27.5</td> <td>467.5</td> </tr> <tr> <td><math>30 &lt; t \leq 35</math></td> <td>24</td> <td>32.5</td> <td>780</td> </tr> <tr> <td><math>35 &lt; t \leq 40</math></td> <td>11</td> <td>37.5</td> <td>412.5</td> </tr> <tr> <td><math>40 &lt; t \leq 45</math></td> <td>9</td> <td>42.5</td> <td>382.5</td> </tr> <tr> <td></td> <td><b>71</b></td> <td></td> <td><b>2267.5</b></td> </tr> </tbody> </table> <p>Estimate for the mean = <math>2267.5 \div 71 = 31.93\dots</math>                      or 32 minutes, to the nearest minute</p>	Time ( <i>t</i> minutes)	Frequency	Midpoint	Frequency × midpoint	$20 < t \leq 25$	10	22.5	225	$25 < t \leq 30$	17	27.5	467.5	$30 < t \leq 35$	24	32.5	780	$35 < t \leq 40$	11	37.5	412.5	$40 < t \leq 45$	9	42.5	382.5		<b>71</b>		<b>2267.5</b>	Multiplying frequencies by your midpoints Dividing the final column total by 71 Correct answer to the nearest minute	1  1  1
Time ( <i>t</i> minutes)	Frequency	Midpoint	Frequency × midpoint																												
$20 < t \leq 25$	10	22.5	225																												
$25 < t \leq 30$	17	27.5	467.5																												
$30 < t \leq 35$	24	32.5	780																												
$35 < t \leq 40$	11	37.5	412.5																												
$40 < t \leq 45$	9	42.5	382.5																												
	<b>71</b>		<b>2267.5</b>																												
29.2 (c)	The actual data values are not known, so it's only possible to estimate the mean finishing time.	Clear explanation	1																												

Question	Answer	Extra information	Marks																
29.3	<p>The midpoint of <math>0 &lt; x \leq 4</math> is 2 The midpoint of <math>4 &lt; x \leq 8</math> is 6</p> <table border="1"> <thead> <tr> <th>Score (x)</th> <th>Frequency</th> <th>Midpoint</th> <th>Frequency × midpoint</th> </tr> </thead> <tbody> <tr> <td><math>0 &lt; x \leq 4</math></td> <td>3y</td> <td>2</td> <td>6y</td> </tr> <tr> <td><math>4 &lt; x \leq 8</math></td> <td>7y</td> <td>6</td> <td>42y</td> </tr> <tr> <td></td> <td><b>10y</b></td> <td></td> <td><b>48y</b></td> </tr> </tbody> </table> <p>Estimate for the mean = <math>48y \div 10y = 4.8</math></p>	Score (x)	Frequency	Midpoint	Frequency × midpoint	$0 < x \leq 4$	3y	2	6y	$4 < x \leq 8$	7y	6	42y		<b>10y</b>		<b>48y</b>	<p>Multiplying frequencies by midpoints Dividing final column total by the frequency total Correct answer of 4.8</p>	<p>1 1 1</p>
Score (x)	Frequency	Midpoint	Frequency × midpoint																
$0 < x \leq 4$	3y	2	6y																
$4 < x \leq 8$	7y	6	42y																
	<b>10y</b>		<b>48y</b>																
29.4	<p>There are 7 data values Lower quartile is the value in position <math>\frac{1}{4}(7+1) = 2</math> This is the missing card Upper quartile is the value in position <math>\frac{3}{4}(7+1) = 6</math> In 6th position, the card value is 8 The IQR = 3, so <math>UQ - LQ = 3</math> <math>8 - LQ = 3</math> Thus, the <math>LQ = 5</math>, which means the missing card's value is 5</p>	<p>Finding the IQR via the LQ and UQ Correct answer</p>	<p>1 1</p>																
29.5 (a)	<p>First arrange the data in ascending order: 63 76 76 78 82 85 87 90 95 99 100 123</p> <p>There are 12 data values LQ will be the mean of the 3rd and 4th values = 77 Median will be the mean of the 6th and 7th values = 86 UQ will be the mean of the 9th and 10th values = 97</p>	<p>1 mark for two of UQ, LQ and median 1 mark for all three</p>	<p>1 1</p>																

Question	Answer	Extra information	Marks
29.5 (b)	It's true that team A have the highest score, but team B has more scores of 100 or more; team A has half of their scores below 86 whereas team B has only two scores less than 86. Team B's median score is also significantly higher than team A's median score. Josh is not correct.	Any two valid observations of the data	2
29.5 (c)	Team A's range is 60 and team B's range is just 35. The IQR for team A is $97 - 77 = 20$ , whereas the IQR for team B is $102 - 88.5 = 13.5$ . Taisa is correct.	Any two valid observations of the data	2
29.6	Either determine the unit cost of grass seed and multiply by the area of the lawn, or, like below, create a ratio relationship: $\frac{\text{£}4.99}{3.66 \text{ m}^2} = \frac{x}{32 \text{ m}^2}$ $x = \frac{32 \times 4.99}{3.66} = \text{£}43.63 \text{ (2 dp)}$ $= \text{£}44 \text{ (to nearest £)}$	Unit cost or ratio established Arrive at the correct multiplication Correct answer	1 1 1

Question	Answer	Extra information	Marks
29.7	<p>If the angles sum to 720, then <math>(n-2) \times 180 = 720</math>, where <math>n</math> is the number of sides. Thus <math>n = 6</math> and the shape is a hexagon.</p> <p>Six sides, with the smallest 20 and the largest 220, and four angles in between them in the sequence. That means there are five “jumps” from 20 to 220.</p> $\frac{220 - 20}{5} = 40$ <p>The common difference is 40, so the six angles are: 20°, 60°, 100°, 140°, 180° and 220°</p>	<p>Find the name / number of sides of the polygon</p> <p>Attempt to identify the arithmetic sequence</p> <p>Fully correct answer</p>	<p>1</p> <p>1</p> <p>1</p>